

Get Help!

We have resources to help you.

For more information or for help to stop using Meth, call:

**GATEWAY: Alcohol and Drug Adult
Managed-Care Services**

1 (800) 488-9919

Crystal Meth Anonymous

Local meetings and additional information

can be found on the website.

www.crystalmeth.org

Narcotics Anonymous

12 step support group meetings in various locations.

(408) 998-4200

For prenatal care information and referrals call

**Santa Clara County Public Health Department
Maternal, Child & Adolescent Health**

1 (800) 310-2332

Produced by

The Coalition for Alcohol & Drug Free Pregnancies

www.cadfp.org

Santa Clara County Public Health Department

Maternal, Child & Adolescent Health

Funding provided by

Santa Clara County Department
of Alcohol & Drug Services (DADS)

If You're Pregnant

Think About Your Baby's Health



Don't Use Meth



If You're Pregnant

Think About Your Baby's Health

Don't Use Meth

Methamphetamine is an illegal and dangerous drug that is highly addictive.

Common names for methamphetamine include meth, crystal, crank, and speed.

If you use meth during your pregnancy, it can hurt your baby's health. Your baby can have problems that can affect the rest of his or her life.

If you have used meth before you knew you were pregnant, stop using meth now. You will give your baby a better chance to be born healthy and without any birth problems that could be lifelong.

If you want to get pregnant, do not use meth. You may not know you are pregnant right away.

The sooner you quit, the healthier your baby will be.

When you use meth during pregnancy, the drug gets into your unborn baby's system and can seriously harm his or her development. It can also affect your baby's brain. Your baby can also be born too early and too small which can lead to many health problems.

Babies born to women who used meth during pregnancy have been described as "irritable babies" and may have:

- Difficulty sucking and swallowing
- Extreme sensitivity to light and human touch
- Shakiness/tremors
- Difficulty sleeping
- Higher heart rate
- Fast or rapid breathing
- Skeletal/bone abnormalities
- A higher chance of dying from Sudden Infant Death Syndrome (SIDS)
- A higher chance of becoming sick from viral infections (such as Hepatitis B and C and HIV).

Some effects of prenatal meth use can be seen years later.

School aged children whose mothers used meth during pregnancy may have:

- Learning difficulties
- Developmental delays
- Negative behavioral patterns
- Irritability
- Hyperactivity
- Attention Deficit Disorder (ADD)
- Fits of anger